



Title	P. gingivalis LPS modulates antimicrobial peptide expression in gingival epithelia
Author(s)	Jin, LJ; Lu, Q; Darveau, R; Samaranayake, LP
Citation	The 84th General Session and Exhibition of the International Association for Dental Research & 1st Meeting of the Pan-Asian-Pacific Federation, Brisbane, Australia, 28 June-1 July 2006. In Journal of Dental Research, 2006, v. 85 Sp Iss B, abstract no. 0898
Issued Date	2006
URL	http://hdl.handle.net/10722/53644
Rights	Creative Commons: Attribution 3.0 Hong Kong License

0898 *P. gingivalis* LPS Modulates Antimicrobial Peptide Expression in Gingival Epithelia

L. JIN¹, Q. LU², R. DARVEAU², and L.P. SAMARANAYAKE¹, ¹The University of Hong Kong, Hong Kong, ²University of Washington, Seattle, USA

Objectives: This study aimed to determine whether *P. gingivalis* LPS could modulate the gingival epithelial expression of human β -defensins (hBDs) through the pattern recognition receptors. **Methods:** Reconstituted human gingival epithelia (RHGE) were incubated with 10 μ g/ml of *P. gingivalis* LPS or 10 ng/ml of *E. coli* LPS for 2, 6, 12, 18, 24, 36, 48, 72 and 96 h. In the blocking assay, the RHGE were pre-incubated with TLR-2 and -4 and CD14 antibodies. The expression of hBDs 1-3, CD14, LBP, TLR-2 and -4, MD-2 and MyD88 mRNAs in RHGE was detected by RT-PCR, while hBDs peptides were detected by immunohistochemistry and ELISA. **Results:** The expression of hBD-2 mRNA was significantly enhanced by *P. gingivalis* LPS in a time-dependant manner within 36 h and remained up to 96 h, which was approximately 3 times higher than that of hBD-1 and -3. Whereas *E. coli* LPS induced a prompt expression of hBD-2 and -3 messages which decreased to basal levels within 36 h. The hBDs peptides were detected in RHGE stimulated with *E. coli* LPS but not in those challenged by *P. gingivalis* LPS. The basal expression of CD14, MD-2 and MyD88 mRNAs was not affected by *P. gingivalis* or *E. coli* LPS. Blockage of CD14, TLR-2 and -4 functions significantly inhibited *E. coli* LPS-induced upregulation of hBD-2 mRNA and peptide expression. However, although combined blockage of TLR-2 and -4 or plus CD14 functions markedly inhibited *P. gingivalis* LPS-induced upregulation of hBD-2 mRNA, it resulted in an increased expression of hBD-2 peptide. **Conclusions:** *P. gingivalis* LPS could enhance the *in vitro* expression of hBDs mRNAs in gingival epithelia likely with the involvement of TLR-2 and -4. While it might evade innate defense surveillance by interference of hBD-2 peptide induction. Supported by Hong Kong Research Grants Council (RGC HKU 7310/00M & 7518/05M).

[Seq #76 - Pathogenesis 2](#)

11:00 AM-12:00 PM, Thursday, 29 June 2006 Brisbane Convention & Exhibition Centre Exhibit Hall 1

[Back to the Periodontal Research - Pathogenesis Program](#)

[Back to the IADR General Session & Exhibition \(June 28 – July 1, 2006\)](#)